## AMENDMENTS TO THE CLAIMS

This listing of claims will replace all prior versions, and listings, of claims in the application:

- 1. (Withdrawn) A method for analyzing at least one of bone mineral density, bone structure and surrounding tissue comprising:
  - a, obtaining an image of a subject;
  - b. locating a region of interest on the image;
  - c. obtaining data from the region of interest; and
- d, deriving quantitative data from the image data obtained at step c, wherein deriving includes extracting data from a measured parameter selected from the group consisting of; a volume of bone marrow edema; a volume of bone marrow edema normalized by at least one of width, area, size, and volume; a volume of osteophytes; a volume of subchondral cysts; a volume of subchondral sclerosis; an area of bone marrow edema; an area of osteophytes; an area of subchondral cysts; an area of subchondral sclerosis; a depth of bone marrow edema; a depth of osteophytes; a depth of subchondral cysts; a depth of subchondral sclerosis; at least one of a volume, area, and depth of at least one of an osteophytes, subchondral cysts, subchondral sclerosis wherein the at least one of volume, area, and depth is normalized by at least one of width, area, size, volume a bone proximal to at least one of the osteophyte, subchondral cyst, or subchondral sclerosis; a volume of menisci; a ratio of volume of normal to at least one of torn, damaged and degenerated meniscal tissue; a ratio of surface area of normal to at least one of torn, damaged and degenerated meniscal tissue; a ratio of surface area of normal to at least one of torn, damaged and degenerated meniscal tissue to total joint or cartilage surface area; a ratio of surface area of at least one of torn, damaged and degenerated meniscal tissue to a total surface area of at least one of joint and cartilage; a size ratio of opposing articular surfaces; a meniscal subluxation/dislocation in millimeters; an index combining different articular parameters; a 3D surface contour information of subchondral bone; an actual or predicted knee flexion angle during gait cycle; a predicted knee rotation during gait cycle; a predicted knee displacement during gait cycle; a predicted load bearing line on cartilage surface during gait cycle and measurement of distance between load bearing line and at least one of cartilage defect and diseased cartilage; a predicted load bearing area on cartilage surface during gait cycle and

measurement of distance between load bearing area and at least one of cartilage defect and

diseased cartilage; a predicted load bearing line on cartilage surface during standing or different degrees of knee flexion and extension and measurement of distance between load bearing line

and at least one of cartilage defect and diseased cartilage; a predicted load bearing area on

cartilage surface during standing or different degrees of knee flexion and extension and

measurement of distance between load bearing area and at least one of cartilage defect and diseased cartilage; a ratio of load bearing area to area of at least one of cartilage defect and

diseased cartilage; a percentage of load bearing area affected by cartilage disease; a location of

cartilage defect within load bearing area; a load applied to cartilage defect, area of diseased

cartilage; and a load applied to cartilage adjacent to at least one of cartilage defect and area of

diseased cartilage.

2. (Withdrawn) The method of claim 1 further including the step of enhancing image data

extracted from the region of interest.

3. (Withdrawn) The method of claim 1 wherein the subject is a mammal.

4. (Withdrawn) The method of claim 1 wherein the subject is a human.

5. (Withdrawn) The method of claim 1 wherein the subject is a horse.

6. (Withdrawn) The method of claim 1 wherein the step of obtaining image data includes

obtaining data from a measured parameter selected from the group consisting of: bone

parameters, cartilage parameters, cartilage defect parameters, cartilage disease parameters, area

parameters, and volume parameters.

7 - 8. (Cancelled)

 $9. \ (Withdrawn) \ A \ method \ for \ analyzing \ at \ least \ one \ of \ bone \ mineral \ density, \ bone \ structure \ and$ 

surrounding tissue comprising:

- a. obtaining an image of a subject;
- b. locating a region of interest on the image;
- c. obtaining data from the region of interest; and
- d. deriving quantitative data from the image data obtained at step c, wherein the image data is obtained of a hip and the step of deriving includes extracting data from a measured parameter selected from the group consisting of: microarchitecture parameters on structures parallel to stress lines; microarchitecture parameters on structures perpendicular to stress lines; geometry; shaft angle; neck angle; diameter of femur neck; hip axis length; largest cross-section of femur head; average thickness of cortical within at least one ROI; standard deviation of cortical thickness within at least one ROI; maximum thickness of cortical within at least one ROI; minimum thickness of cortical within at least one ROI; minimum thickness of cortical within at least one ROI; and hip joint space width.
- 10. (Withdrawn) A method for analyzing at least one of bone mineral density, bone structure and surrounding tissue comprising:
  - a. obtaining an image of a subject;
  - b. locating a region of interest on the image;
  - c. obtaining data from the region of interest; and
- d. deriving quantitative data from the image data obtained at step c, wherein the image data is obtained from a region of a spine and the step of deriving includes extracting data from a measured parameter selected from the group consisting of: microarchitecture parameters on vertical structures; microarchitecture parameters on horizontal structures; geometry; superior endplate cortical thickness; inferior endplate cortical thickness; anterior vertebral wall cortical thickness; posterior vertebral wall cortical thickness; superior aspect of pedicle cortical thickness; inferior aspect of pedicle cortical thickness; vertebral height; vertebral diameter; pedicle thickness, maximum vertebral height; minimum vertebral height; average vertebral height; anterior vertebral height; medial vertebral height; posterior vertebral height; maximum inter-vertebral height; minimum inter-vertebral height; and average inter- vertebral height.

## 11. (Cancelled)

- 12. (Currently Amended) A method for analyzing bone, comprising:
  - a. obtaining image data of an anatomical structure;
  - b. defininglocating a region of interest in the image data that includes trabecular bone;
- c. defining within the region of interest at least first and second windowsgenerating a data structure representing a trabecular pattern within the region of interest:
- d. analyzing information in the first window to determine at least a first value for a parameter selected from the group comprising bone microarchitecture; bone macroanatomy; cartilage microarchitecture; cartilage macroanatomy; connective tissue microarchitecture; connective tissue macroanatomy; or biomechanical parametersperforming pixel-neighborhood analysis on the at least one data structure; and
- e. analyzing information in the second window to determine at least a second value for the parameter; and

f. generating a set of data at least in part from the first and second values deriving quantitative data from the results of the pixel neighborhood analysis, wherein the step of deriving includes combining at least two parameters from the group consisting of: total area; trabecular perimeter; node count (N); segment count (S); and node-to-free-end segment count (NF).

- 13. (Currently Amended) The method of claim [[173]]12 wherein the first and second values for the parameter are determined using pixel neighborhood analysistrabeeular bone pattern factor is (P1-P2)/(A1-A2) further wherein P1 and A1 are the perimeter length and trabeeular bone area before dilation and P2 and A2 corresponding values after a single pixel dilation.
- 14. (Withdrawn) A method for analyzing at least one of bone mineral density, bone structure and surrounding tissue comprising:
  - a. obtaining an image of a subject;
  - b. locating a region of interest on the image;
  - c. obtaining data from the region of interest; and
- d. deriving quantitative data from the image data obtained at step c, wherein the step of deriving includes extracting at least one of cartilage parameters, cartilage defect parameters, and diseased cartilage parameters, wherein the data extracted is selected from the group consisting

of: mean cartilage thickness over substantially total surface; mean cartilage thickness in focal area; maximum cartilage thickness in focal area; minimum cartilage thickness in focal area; 3D cartilage shape information; cartilage curvature analysis; volume of cartilage defect/diseased cartilage; depth of cartilage defect/diseased cartilage; area of cartilage defect/diseased cartilage; at least one of 2D and 3D location of cartilage defect/diseased cartilage in the articular surface; at least one of 2D and 3D location of cartilage defect/diseased cartilage in relationship to weight-bearing area; a ratio of at least two of diameter of cartilage defect, diameter of diseased cartilage, and thickness of surrounding normal cartilage; a ratio of at least two of depth of cartilage defect, depth of diseased cartilage and thickness of surrounding normal cartilage; a ratio of at least two of volume of cartilage defect, volume of diseased cartilage and thickness of surrounding normal

15. (Withdrawn) The method of claim 1 wherein at least one step is performed automatically.

cartilage; a ratio of at least two of surface area of cartilage defect, surface area of diseased cartilage and total joint surface area; and a ratio of at least two of volume of cartilage defect,

16. (Withdrawn) The method of claim 1 wherein at least one step is performed semiautomatically.

volume of diseased cartilage and total cartilage volume.

- 17. (Withdrawn) The method of claim 1 wherein at least one of the steps is performed on a first computer.
- 18. (Withdrawn) The method of claim 1 wherein at least one of the steps is performed on a first computer and at least one of the steps is performed on a second computer.
- 19. (Withdrawn) The method of claim 18 wherein the first computer and the second computer are connected by one of a peer to peer network, direct link, intranet, and internet.
- 20. (Withdrawn) The method of claim 1 wherein the step of locating a region of interest is repeated.

21. (Withdrawn) The method of claim 1 wherein the step of obtaining image data from a region of interest is repeated.

22. (Withdrawn) The method of claim 1 wherein at least one of the image and data is converted to a 2D pattern.

23. (Withdrawn) The method of claim 22 wherein the 2D pattern is evaluated.

24. (Withdrawn) The method of claim 22 wherein the 2D pattern is converted to a 3D pattern.

25. (Withdrawn) The method of claim 23 wherein the 2D pattern is converted to a 3D pattern.

26. (Withdrawn) The method of claim 22 wherein the 2D pattern is converted to a 4D pattern.

27. (Withdrawn) The method of claim 23 wherein the 2D pattern is converted to a 4D pattern.

28. (Withdrawn) The method of claim 25 wherein the 3D pattern is converted to a 4D pattern.

29. (Withdrawn) The method of claim 1 wherein at least one of the image and data is converted to a 3D pattern.

30. (Withdrawn) The method of claim 29 wherein the 3D pattern is evaluated.

31. (Withdrawn) The method of claim 29 wherein the 3D pattern is converted to a 2D pattern.

32. (Withdrawn) The method of claim 31 wherein the 3D pattern is converted to a 2D pattern.

33. (Withdrawn) The method of claim 29 wherein the 3D pattern is converted to a 4D pattern.

Reply to Office Action dated December 28, 2010

34. (Withdrawn) The method of claim 30 wherein the 3D pattern is converted to a 4D pattern.

35. (Withdrawn) The method of claim 31 wherein the 2D pattern is converted to a 4D pattern.

36. (Withdrawn) The method of claim 1 wherein the at least one of the image and data is

converted to a 4D pattern.

37. (Withdrawn) The method of claim 36 wherein the 4D pattern is evaluated.

38. (Withdrawn) The method of claim 1 further comprising the step of administering a candidate

agent.

39. (Withdrawn) The method of claim 38 wherein the candidate agent is at least one agent

selected from the group consisting of: substance administered to a subject, substance ingested by

 $a\ subject,\ molecules,\ pharmaceuticals,\ biopharmaceuticals,\ agropharmaceuticals.$ 

40. (Withdrawn) The method of claim 1 further comprising the step of comparing the data

obtained to a database.

41. (Withdrawn) The method of claim 1 further comprising the step of comparing the data

obtained to a subset of a database.

42. (Withdrawn) The method of claim 1 further comprising the step of comparing the

quantitative data to an image taken at T1.

43. (Withdrawn) The method of claim 1 further comprising the step of comparing the

quantitative data to an image taken prior to the image under analysis.

44. (Withdrawn) The method of claim 1 further comprising the step of comparing the

quantitative data to an image taken at Tn.

45. (Withdrawn) The method of claim 1 further comprising the step of transmitting at least one of the image and data.

46. (Withdrawn) The method of claim 45 further comprising the step of analyzing the converted image.

47. (Withdrawn) The method of claim 46 wherein the step of analyzing the converted image occurs at least at one of prior to transmitting the image or after transmitting the image.

48. (Withdrawn) The method of claim 1 wherein the image is converted to at least one of a pattern of normal, diseased, and normal and diseased.

49. (Withdrawn) The method of claim 1 wherein obtaining image data includes measuring at least one of microarchitecture and macroanatomical structures.

50. (Withdrawn) The method of claim 49 further comprising measuring the average density.

51. (Withdrawn) The method of claim 50 wherein the average density measurement includes a calibrated density of the region of interest.

52. (Withdrawn) The method of claim 49 further comprising measuring microanatomical structures on at least one of dental, spine, hip, knee and bone core x-rays.

53. (Withdrawn) A method for analyzing at least one of bone mineral density, bone structure and surrounding tissue comprising:

a. obtaining an image of a subject;

b. locating a region of interest on the image;

c. obtaining data from the region of interest; and

Appl. No. 10/665,725 Amendment dated June 28, 2011

Reply to Office Action dated December 28, 2010

d. deriving quantitative data from the image data obtained at step c, wherein the step of

deriving includes measuring at least one of: calibrated density of extracted structures; calibrated density of background; average intensity of extracted structures; average intensity of background

area wherein the background area includes non-extracted structures; structural contract wherein

structural contrast is an average intensity of extracted structures divided by an average intensity

of a background; calibrated structural contrast wherein calibrated structural contrast is a

calibrated density of extracted structures divided by a calibrated density of a background; total

area of extracted structures; total area of a region of interest; an area of extracted structures normalized by a total area of a region of interest; a boundary length of an extracted area

normalized by a total area of a region of interest; a number of structures normalized by an area of

a region of interest; a trabecular bone pattern factor; a measurement of concavity and convexity

a region of interest, a tracecular cone partern factor, a measurement of concurry and conversity

of structures; a star volume of extracted structures; a star volume of background; and a number

of loops normalized by an area of a region of interest.

54. (Withdrawn) The method of claim 49 further comprising measuring a distance transform of

extracted structures.

55. (Withdrawn) The method of claim 54 wherein the measurement on the distance transform of

extracted structures further comprises one or more of: an average regional maximum thickness; a

standard deviation of regional maximum thickness; a largest value of regional maximum  $\,$ 

thickness; and a median regional maximum thickness.

56 - 88 (Cancelled)

89. (Previously Presented) The method of claim 12 further including the step of enhancing the

image data.

90. (Currently Amended) The method of claim 12 wherein the image datasubject is of a

mammal.

91. (Currently Amended) The method of claim 12 wherein the image datasubject is of a human.

92. (Currently Amended) The method of claim 12, further comprising generating a field of

sampling points within the region of interest, the first window being located at a first sampling point and the second window being located at a second sampling point-wherein the subject is a

horse.

93. (Previously Presented) The method of claim 12 wherein at least one step is performed

automatically.

94. (Previously Presented) The method of claim 12 wherein at least one step is performed semi-

automatically.

95. (Previously Presented) The method of claim 12 wherein at least one of the steps is performed

on a first computer.

96. (Previously Presented) The method of claim 12 wherein at least one of the steps is performed

on a first computer and at least one of the steps is performed on a second computer.

97. (Previously Presented) The method of claim 96 wherein the first computer and the second

computer are connected by one of a peer to peer network, direct link, intranet, and internet.

98. (Currently Amended) The method of claim 12 wherein the image is an x-ray imagestep of

locating a region of interest is repeated.

99. (Currently Amended) The method of claim 12 wherein the <u>first and second windows abut</u>

without overlappingstep of extracting a trabecular pattern from a region of interest is repeated.

100. (Cancelled)

101. (Currently Amended) The method of claim 12 wherein the first and second windows do not overlaptrabecular pattern is a 2D pattern.

102. (Currently Amended) The method of claim 12 wherein the image data is oftrabecular pattern is converted to a 3D imagepattern.

103. (Currently Amended) The method of claim [101]92 wherein the step of generating a field of sampling points within the region of interest comprises generating an irregularly spaced field of sampling points within the region of interest2D pattern is converted to a 3D pattern.

104. (Currently Amended) The method of claim [12]92 wherein the step of generating a field of sampling points within the region of interest comprises generating a regularly spaced field of sampling points within the region of interest trabecular pattern is converted to a 4D pattern.

105. (Currently Amended) The method of claim [101]12 wherein the bone is osteoporotic 2D pattern is converted to a 4D pattern.

106. (Withdrawn) The method of claim 103 wherein the 3D pattern is converted to a 4D pattern.

107. (Withdrawn) The method of claim 12 wherein at least one of the image and data is converted to a 3D pattern.

108. (Withdrawn) The method of claim 107 wherein the 3D pattern is evaluated.

109. (Withdrawn) The method of claim 107 wherein the 3D pattern is converted to a 2D pattern.

110. (Withdrawn) The method of claim 108 wherein the 3D pattern is converted to a 2D pattern.

111. (Withdrawn) The method of claim 107 wherein the 3D pattern is converted to a 4D pattern.

Appl. No. 10/665,725 Amendment dated June 28, 2011

Reply to Office Action dated December 28, 2010

112. (Withdrawn) The method of claim 108 wherein the 3D pattern is converted to a 4D pattern.

113. (Withdrawn) The method of claim 109 wherein the 2D pattern is converted to a 4D pattern.

114. (Withdrawn) The method of claim 12 wherein the at least one of the image and data is

converted to a 4D pattern.

115. (Withdrawn) The method of claim 114 wherein the 4D pattern is evaluated.

116. (Previously Presented) The method of claim 12 further comprising the step of administering

a candidate agent.

117. (Previously Presented) The method of claim 116 wherein the candidate agent is at least one

agent selected from the group consisting of: substance administered to a subject, substance

ingested by a subject, molecules, pharmaceuticals, biopharmaceuticals, agropharmaceuticals.

118. (Currently Amended) The method of claim 12 further comprising the step of comparing the

set of generated data-obtained to a database.

119. (Currently Amended) The method of claim 12 further comprising the step of comparing the

set of data generatedobtained to a subset of a database.

120. (Currently Amended) The method of claim [12]117 further comprising the step of

comparing the set of quantitative data to [[an]]a corresponding set of data generated from an

image taken prior to administration of the candidate agent at T1.

121. (Currently Amended) The method of claim 12 further comprising the step of comparing the

set ofquantitative data to [[an]]a corresponding set of data generated from an image taken prior

to the image under analysis.

Appl. No. 10/665,725 Amendment dated June 28, 2011

Reply to Office Action dated December 28, 2010

122. (Currently Amended) The method of claim 12, wherein the image data is twodimensional further comprising the step of comparing the quantitative data to an image taken at

Tn.

123. (Currently Amended) The method of claim 12, wherein the image data is four-

dimensional further comprising the step of transmitting at least one of the image data and

quantitative data.

124. (Currently Amended) The method of claim [[123]]12, further comprising creating a

parameter map based at least in part on the values associated with the parameter wherein the

image data is converted to at least one of a pattern of normal, diseased, and normal and diseased.

125. (Currently Amended) The method of claim [[124]]12, wherein the first and second windows

are of different resolution further comprising the step of analyzing the converted image data.

126. (Currently Amended) The method of claim [[125]]12 wherein the first and second windows

are of different sizes step of analyzing the converted image occurs at least at one of prior to transmitting the image or after transmitting the image data.

127. Cancelled.

128. (Currently Amended) The method of claim 12, wherein the first and second windows have

different orientations further comprising measuring the average density.

129. (Currently Amended) The method of claim 128 wherein the first and second windows are of

different sizes average density measurement includes a calibrated density of the region of interest.

130. Cancelled.

131. (Withdrawn) The method of claim 14 further including the step of enhancing image data extracted from the region of interest.

132. (Withdrawn) The method of claim 14 wherein the subject is a mammal.

133. (Withdrawn) The method of claim 14 wherein the subject is a human.

134. (Withdrawn) The method of claim 14 wherein the subject is a horse.

135. (Withdrawn) The method of claim 14 wherein at least one step is performed automatically.

136. (Withdrawn) The method of claim 14 wherein at least one step is performed semiautomatically.

137. (Withdrawn) The method of claim 14 wherein at least one of the steps is performed on a first computer.

138. (Withdrawn) The method of claim 14 wherein at least one of the steps is performed on a first computer and at least one of the steps is performed on a second computer.

139. (Withdrawn) The method of claim 138 wherein the first computer and the second computer are connected by one of a peer to peer network, direct link, intranet, and internet.

140. (Withdrawn) The method of claim 14 wherein the step of locating a region of interest is repeated.

141. (Withdrawn) The method of claim 14 wherein the step of obtaining image data from a region of interest is repeated. 142. (Withdrawn) The method of claim 14 wherein at least one of the image and data is converted to a 2D pattern.

143. (Withdrawn) The method of claim 142 wherein the 2D pattern is evaluated.

144. (Withdrawn) The method of claim 142 wherein the 2D pattern is converted to a 3D pattern.

145. (Withdrawn) The method of claim 143 wherein the 2D pattern is converted to a 3D pattern.

146. (Withdrawn) The method of claim 142 wherein the 2D pattern is converted to a 4D pattern.

147. (Withdrawn) The method of claim 143 wherein the 2D pattern is converted to a 4D pattern.

148. (Withdrawn) The method of claim 145 wherein the 3D pattern is converted to a 4D pattern.

149. (Withdrawn) The method of claim 14 wherein at least one of the image and data is converted to a 3D pattern.

150. (Withdrawn) The method of claim 149 wherein the 3D pattern is evaluated.

151. (Withdrawn) The method of claim 149 wherein the 3D pattern is converted to a 2D pattern.

152. (Withdrawn) The method of claim 150 wherein the 3D pattern is converted to a 2D pattern.

153. (Withdrawn) The method of claim 149 wherein the 3D pattern is converted to a 4D pattern.

154. (Withdrawn) The method of claim 150 wherein the 3D pattern is converted to a 4D pattern.

155. (Withdrawn) The method of claim 151 wherein the 2D pattern is converted to a 4D pattern.

156. (Withdrawn) The method of claim 14 wherein the at least one of the image and data is converted to a 4D pattern.

157. (Withdrawn) The method of claim 156 wherein the 4D pattern is evaluated.

158. (Withdrawn) The method of claim 14 further comprising the step of administering a candidate agent.

159. (Withdrawn) The method of claim 158 wherein the candidate agent is at least one agent selected from the group consisting of: substance administered to a subject, substance ingested by a subject, molecules, pharmaceuticals, biopharmaceuticals, agropharmaceuticals.

160. (Withdrawn) The method of claim 14 further comprising the step of comparing the data obtained to a database.

161. (Withdrawn) The method of claim 14 further comprising the step of comparing the data obtained to a subset of a database.

162. (Withdrawn) The method of claim 14 further comprising the step of comparing the quantitative data to an image taken at T1.

163. (Withdrawn) The method of claim 14 further comprising the step of comparing the quantitative data to an image taken prior to the image under analysis.

164. (Withdrawn) The method of claim 14 further comprising the step of comparing the quantitative data to an image taken at Tn.

165. (Withdrawn) The method of claim 14 further comprising the step of transmitting at least one of the image and data.

166. (Withdrawn) The method of claim 165 further comprising the step of analyzing the converted image.

167. (Withdrawn) The method of claim 166 wherein the step of analyzing the converted image occurs at least at one of prior to transmitting the image or after transmitting the image.

168. (Withdrawn) The method of claim 14 wherein the image is converted to at least one of a pattern of normal, diseased, and normal and diseased.

169. (Withdrawn) The method of claim 14 wherein obtaining image data includes measuring at least one of microarchitecture and macroanatomical structures.

170. (Withdrawn) The method of claim 169 further comprising measuring the average density.

171. (Withdrawn) The method of claim 170 wherein the average density measurement includes a calibrated density of the region of interest.

172. (Withdrawn) The method of claim 171 further comprising measuring microanatomical structures on at least one of dental, spine, hip, knee and bone core x-rays.

173. (Currently Amended) The method of claim [[12]]]3 wherein the parameter comprises bone microarchitecturequantitative data includes an additional parameter-from the group of parameters consisting of: trabecular contrast wherein the trabecular contrast is determined as one of the trabecular equivalent thickness and marrow equivalent thickness; trabecular area; marrow distance transform; marrow distance transform regional maxima values; connected skeleton count or trees (T); node-to-node segment count (NN); node-to-node segment length (NNL) node-to-free-end segment length (NNL); free-end-to-free-end segment length (FF.TSL); total struts length (TSL); FF.TSL/TSL; NN.TSL/TSL; loop count (Lo); loop area; mean distance transform values for each segment (Tb.Th); mean distance transform values for each node-to-node segment

(Tb.Th.NN); mean distance transform values for each node-to-free-end segment (Tb.Th.NF); orientation of each segment; angle of each segment; angle between segments; length-thickness ratios (NNL/Tb.Th.NN) and (NFL/Tb.Th.NF); and interconnectivity index (ICI) ICI=(N\*NN)/(T\*(NF+1)).

- 174. (Withdrawn) A method for evaluating musculoskeletal tissue, comprising:
  - a. obtaining image data representing tissue;
  - b. defining a region of interest in the tissue that includes bone tissue;
  - c. extracting trabecular bone structure from the region of interest;
- d. processing the trabecular bone structure to obtain data representing the bone structure; and
  - e. measuring a parameter based on the data representing the trabecular bone structure.
- 175. (Withdrawn) The method of claim 174, wherein extracting further comprises amplifying image data.
- 176. (Withdrawn) The method of claim 174, wherein extracting further comprises background subtraction and thresholding of image data.
- 177. (Withdrawn) The method of claim 174, wherein extracting further comprises edge filtering and thresholding of image data.
- 178. (Withdrawn) The method of claim 174, wherein processing the trabecular bone structure further comprises performing a distance transform.
- 179. (Withdrawn) The method of claim 174, wherein processing the trabecular bone structure further comprises skeletonizing the extracted trabecular bone structure.
- 180. (Withdrawn) The method of claim 174, wherein processing the trabecular bone structure data further comprises segmenting the extracted trabecular bone structure.

181. (Withdrawn) The method of claim 174, wherein processing the trabecular bone structure further comprises performing watershed segmentation.

182. (Withdrawn) The method of claim 174, wherein the parameter is from a group of parameters consisting of; trabecular contrast wherein the trabecular contrast is determined as one of the trabecular equivalent thickness and marrow equivalent thickness; at least one of trabecular area and total area; trabecular perimeter; trabecular distance transform; marrow distance transform; trabecular distance transform regional maxima values; marrow distance transform regional maxima values; star volume; trabecular bone pattern factor; connected skeleton count or trees (T); node count (N); segment count (S); node-to-node segment count (NN); node-to-freeend segment count (NF); node-to-node segment length (NNL) node-to-free-end segment length (NFL); free-end-to-free-end segment length (FFL); node-to-node total struts length (NN.TSL); free-end-to-free-ends total struts length (FF,TSL); total struts length (TSL); FF,TSL/TSL; NN.TSL/TSL; loop count (Lo); loop area; mean distance transform values for each connected skeleton; mean distance transform values for each segment (Tb.Th); mean distance transform values for each node-to-node segment (Tb.Th.NN); mean distance transform values for each node-to-free-end segment (Tb.Th.NF); orientation of each segment; angle of each segment; angle between segments; length-thickness ratios (NNL/Tb.Th.NN) and (NFL/Tb.Th.NF); and interconnectivity index (ICI) ICI=(N\*NN)/(T\*(NF+1)).

- 183. (Withdrawn) A method for evaluating bone tissue, comprising:
  - a. obtaining image data representing an anatomical structure that includes bone tissue;
  - b. extracting a trabecular bone pattern from at least a portion of the image data;
  - c. skeletonizing the trabecular bone pattern; and
  - d. quantifying data based on the skeletonized trabecular bone pattern.

184. (Withdrawn) A method for evaluating trabecular bone tissue, comprising quantifying one or more parameters at least in part from a skeleton data structure of a trabecular bone pattern extracted from an image including the trabecular bone. Reply to Office Action dated December 28, 2010

185. (Withdrawn) A method for analyzing bone using a computer system, comprising:

- a. obtaining image data of an anatomical structure;
- b. locating a region of interest in the image data that includes trabecular bone;
- c. determining a trabecular pattern from the region of interest;
- d. determining a cortical bone parameter of the anatomical structure;
- f. deriving quantitative data based at least in part on the cortical bone parameter and the data structure.
- 186. (Withdrawn) The method of claim 185 further including the step of enhancing the image data.
- 187. (Withdrawn) The method of claim 185 wherein the cortical bone parameter is determined from the region of interest.
- 188. (Withdrawn) The method of claim 185 wherein the subject is a human.
- 189. (Withdrawn) The method of claim 185 wherein at least one step is performed automatically.
- 190. (Withdrawn) The method of claim 185 wherein at least one step is performed semiautomatically.
- 191. (Withdrawn) The method of claim 185 wherein the trabecular pattern is a 2D pattern.
- 192. (Withdrawn) The method of claim 191 wherein the trabecular pattern is converted to a 3D pattern.
- 193. (Withdrawn) The method of claim 191 wherein the trabecular pattern is converted to a 4D pattern.

Reply to Office Action dated December 28, 2010

194. (Withdrawn) The method of claim 185 further comprising the step of administering a candidate agent.

195. (Withdrawn) The method of claim 185 further comprising the step of saving the data obtained to a database

196. (Withdrawn) The method of claim 185 wherein obtaining image data includes obtaining image data of at least one of microarchitecture and macroanatomical structures.

197. (Withdrawn) The method of claim 185 further comprising measuring microanatomical structures on at least one of dental, spine, hip, knee and bone core x-rays.

198. (Withdrawn) A method for analyzing bone using a computer system, comprising:

a, obtaining image data of an anatomical structure that includes cortical and trabecular bone:

- b. locating a region of interest in the image data that includes trabecular bone;
- c. determining a trabecular bone pattern from the region of interest;
- d. determining a thickness of cortical bone of the anatomical structure;
- f. deriving quantitative data based at least in part on the thickness and the trabecular bone pattern.

199. (Withdrawn) The method of claim 198 further including the step of enhancing the image data.

200. (Withdrawn) The method of claim 198 wherein the thickness of cortical bone is determined from the region of interest.

201. (Withdrawn) The method of claim 198 wherein the subject is a human.

202. (Withdrawn) The method of claim 198 wherein at least one step is performed automatically.

203. (Withdrawn) The method of claim 198 wherein at least one step is performed semiautomatically.

204. (Withdrawn) The method of claim 198 wherein the trabecular pattern is a 2D pattern.

205. (Withdrawn) The method of claim 204 wherein the trabecular pattern is converted to a 3D pattern.

206. (Withdrawn) The method of claim 204 wherein the trabecular pattern is converted to a 4D pattern.

207. (Withdrawn) The method of claim 198 further comprising the step of administering a candidate agent.

208. (Withdrawn) The method of claim 198 further comprising the step of saving the data obtained to a database.

209. (Withdrawn) The method of claim 198 wherein obtaining image data includes obtaining image data of at least one of microarchitecture and macroanatomical structures.

210. (Withdrawn) The method of claim 198 further comprising measuring microanatomical structures on at least one of dental, spine, hip, knee and bone core x-rays.